



Influence of Diet on the Effect of the Probiotic *Lactobacillus paracasei* in Rats Suffering From Allergic Asthma

Ao Xie, Jiaping Song, Shan Lu, Yinhui Liu, Li Tang and Shu Wen*

Department of Microecology, College of Basic Medical Sciences, Dalian Medical University, Dalian, China

OPEN ACCESS

Edited by:

Tingtao Chen,
Nanchang University, China

Reviewed by:

Hong Wei,
Army Medical University, China
Jiaming Liu,
Wenzhou Medical University, China
Longxian Lv,
Zhejiang University, China

*Correspondence:

Shu Wen
shuwen@dmu.edu.cn

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Microbiology

Received: 07 July 2021

Accepted: 31 August 2021

Published: 27 September 2021

Citation:

Xie A, Song J, Lu S, Liu Y, Tang L
and Wen S (2021) Influence of Diet on
the Effect of the Probiotic
Lactobacillus paracasei in Rats
Suffering From Allergic Asthma.
Front. Microbiol. 12:737622.
doi: 10.3389/fmicb.2021.737622

Mounting evidence suggests that probiotics can be used to treat allergic asthma by modulating the gut microbiota, and that the effects of probiotics may be influenced by environmental factors such as diet. We conducted a rat model with allergic asthma (AA) modulated by *Lactobacillus paracasei*, feeding up with high-fat or high-fiber diets based on collecting data from 85 questionnaires. The systemic proinflammatory cytokines were detected by ELISA and the overall structure of fecal microbiota was analyzed via 16S rRNA gene sequencing. The results showed consumption of a high-fiber diet alleviated the allergic symptoms and airway inflammation, and led to improving the imbalance of T-helper type 1 (Th1)/Th2 cells with increased expression of interferon- γ and decreased expression of interleukin-4. Whereas, the high-fat diet had deteriorating implications and skewed the inflammatory perturbation. Furthermore, abundances of phylum Bacteroidetes, families Muribaculaceae, Tannerellaceae, Prevotellaceae, Enterococcaceae, genera Allobaculum, Parabacteroides, and Enterococcus were enriched in *L. paracasei*-modulating rats fed with high-fiber diet. Firmicutes and Proteobacteria, families Lachnospiraceae, Ruminococcaceae and Desulfovibrionaceae, genera Blautia, unidentified_Ruminococcaceae, unidentified_Clostridiales and Oscillibacter were in relatively high abundance in the rats administered high-fat diet. Association between changed microbiota and inflammatory cytokines was also conferred. These data indicated that the efficacy of *L. paracasei* in allergic asthma was influenced by different dietary patterns. Hence, diet is important for probiotic therapy when managing allergic asthma.

Keywords: diet pattern, intestinal microbiota, *Lactobacillus paracasei*, allergic asthma, airway inflammation

INTRODUCTION

There has been an increase in the prevalence of allergic asthma (AA) globally, particularly in children and in Western countries, which has led to substantial financial and medical burdens (Colombo et al., 2019). The recommended resolution for AA is to use high-dose inhaled corticosteroids (ICs) alone or combine them with long-acting bronchodilators. Unfortunately, besides the undesirable effects of long-term treatment with ICs, a significant number of asthmatic patients fail to respond to IC therapy.

A disturbed balance of Th1/Th2 cells and Th17/T-regulatory cells (T_{regs}) imbalance in AA have been reported. Th2 cytokines [e.g., interleukin(IL)-4, IL-5, and IL-13] and potent proinflammatory cytokines (e.g., IL-1 β , IL-6, IL-17, IL-25, and tumor necrosis factor (TNF)- α) orchestrate mucosal inflammation. Inflammatory mediators have been shown to impair or boost mucosal inflammation in the airways. Over recent decades, increased perturbation of the microbiota has been demonstrated to contribute to the development of allergic inflammation. Mounting evidence suggests that low diversity of the gut microbiota (GM) in early infancy is a vital risk factor for the development of immune-mediated allergic diseases (Abrahamsson et al., 2014; Tsabouri et al., 2014; Dzidic et al., 2017). Furthermore, microbial colonization in germ-free mice within the first days of life or in antibiotic-treated mice has been indicated to protect against increased IgE levels (Herbst et al., 2011) and promote tolerance to aeroallergens via T_{regs} induction (Gollwitzer et al., 2014).

Beneficial bacteria modulate the intestinal microbiota and mucosal immune responses. A significantly decreased profile of Th2 cytokines in response to probiotic treatment has been found in animal experiments (Choi et al., 2018; Makino et al., 2019). Oral administration of mixed strains of *Clostridium* species to BALB/c mice has been shown to stimulate the allergen-induced expansion of T_{regs} in the colonic mucosa and decreased systemic production of IgE (Atarashi et al., 2011). *Lactobacillus paracasei*, commonly used in dairy-product fermentation, has been shown to mitigate respiratory tract allergies in various studies applying in mice and humans (Fujiwara et al., 2004; Schabussova et al., 2012; Wang et al., 2017; Lin et al., 2020). However, clinical trials have conflicting results in assessing *L. paracasei* prevents allergy (Kuitunen et al., 2009; Fiocchi et al., 2012; Jensen et al., 2012; West et al., 2013).

Several studies have underlined the critical role of diet in AA pathogenesis. High consumption of fat (e.g., Western diet) is likely related to an increased risk of asthma (Hou et al., 2011; Protudjer et al., 2012; Manzel et al., 2014; Myles, 2014; Thorburn et al., 2014; Julia et al., 2015). Diets high in fiber (e.g., fruit and vegetables) protect from allergies and are associated with decreased inflammation in the airways (Ellwood et al., 2001; Maslowski et al., 2009; Frei et al., 2012; Grimshaw et al., 2014; Trompette et al., 2014). The effect of diets on allergic asthma might have a close association with the intestinal microbiota. Different diets yield numerous metabolites that can influence the immune response, such as short-chain fatty acids (SCFAs) fermented by colonic commensal, responsible for the effect of anti-inflammation, and the saturated fatty acids or cholesterol displayed exacerbation for allergic disease. However, how to promote the effect of probiotics in people with different diets is not known.

Herein, we collected data (via questionnaires) from 85 human participants to design specific different dietary patterns and established a rat model of AA (using ovalbumin(OVA)) fed up with the designed diets to assess the efficacy of different diets on *L. paracasei*-modulated allergic inflammation. Our findings emphasize the importance of diet to explain the effect of *L. paracasei* on AA prevention.

MATERIALS AND METHODS

Collection of Dietary Data

Collection of dietary data was conducted from 25 November to 2 December 2017. The food inventory was documented according to the Chinese Dietary Guidelines (fourth edition, 2016) and AA history. Food (including edible oils and salt) consumption was determined by weighing, individuals aged >18 years, and excluding those unable to complete data collection.

Preparation of Probiotics


Lactobacillus paracasei DMLA16017 (Department of Microecology, Dalian Medical University) was grown in MRS broth (Hopebio, Qingdao, China) at pH 5.7 \pm 0.2 and under anaerobic conditions. Bacteria were harvested in the early stationary phase, washed with phosphate-buffered saline (PBS; Solarbio, Beijing, China), and stored at -80°C . Cell counts were determined by plating serial dilutions.

Animals and Intervention Design

The study protocol was approved by the Animal Ethics Committee of Dalian Medical University (No. L201312). Specific pathogen-free (SPF) three-week-old male Wistar rats were purchased from the Animal Center of Dalian Medical University and allowed to acclimatize to their surroundings for ≥ 24 h before experimentation. Rats were housed under standard temperature conditions and humidity with a 12-h light-dark schedule with free access to sterile food and autoclaved water.

An AA model was established according to a modification of airway inflammation described by Zhang and Shi (Zhang et al., 2016). After acclimatization, rats were assigned randomly to five groups ($n = 8/\text{group}$): the control group (Con), the model group (Model), the probiotics-modulated group (Probio), the high-fat diet intervened probiotics-modulated group (Hfat), and the high-fiber diet intervened probiotics-modulated group (Hfiber). Sensitizations were undertaken on days 1, 5, 8, and 12 in all groups except the Con group. Rats were sensitized to OVA (egg-white albumin of chickens $\geq 98\%$; Sigma-Aldrich, Saint Louis, MO, United States) by intraperitoneal injection of 0.2 mL of alum-precipitated antigen [comprising 10 μg of OVA absorbed into 1 mg of aluminum hydroxide (Sigma-Aldrich)]. From day 15, rats were administered (i.n.) 20 μL of 1% OVA diluted in PBS for 7 consecutive days and exposed daily to 1% OVA aerosol in PBS in an aerosol cabin for 30 min between day 28 and day 32 in the challenge phase. Rats in the Con group received PBS alone (Figure 1).

After developing airway inflammation, other groups except Con and Model group were gavage with 1.0×10^9 colony-forming units of *L. paracasei* suspended in 0.2 mL of PBS per day, 1 h before the challenge. Rats that underwent gavage with *L. paracasei* were received high-fat (Hfat), high-fiber (Hfiber), or basic chow (Con, Model and Probio), respectively, that we purchased from Huafukang Bioscience (Beijing, China) (see Supplementary Table 1 for nutritional parameters). The intervention was from day 16 until day 32 (Figure 1). Twenty-four hours after the final challenge, all animals were sacrificed.



Evaluation of Allergy Symptoms

For each rat, the frequency of two symptoms of allergy (nose rubbing and sneezing) was recorded immediately for 15 min after the last challenge with OVA aerosol on day 32 in a blinded manner, as described previously (Shinmei et al., 2009).

Cell Counts for Bronchoalveolar Lavage Fluid (BALF)

After sacrifice on day 33, lungs were perfused with 1 mL of PBS containing 10% fetal bovine serum through a tracheal cannula to collect BALF. The latter was centrifuged at $1000 \times g$ for 10 min at room temperature, and the supernatant was separated and stored until use. The pellet was resuspended in PBS for cell counting. BALF cells contain leukocytes and provide a “snapshot” of airway disease.

Enzyme-Linked Immunosorbent Assay (ELISA) for Total Level of OVA-Specific IgE and Cytokines

Twenty-four hours after the final challenge with OVA, rats were anesthetized and blood samples were withdrawn by cardiac puncture. Serum was separated by centrifugation ($1000 \times g$, 10 min, room temperature) after resting for 1 h. The OVA-specific IgE level in serum was determined with an ELISA kit (No. RX-D302482R) according to manufacturer (Ruixinbio, Quanzhou, China) instructions. Absorbance was measured at 450 nm.

The level of IL-4, IL-17, interferon(IFN)- γ , and TGF- β in BALF and serum prepared as stated above were measured by ELISA kits (No. SEA077Ra, No. SEA063Ra, No. SEA049Ra; and No. RX302048R) following manufacturer (Cloud-Clone, Wuhan, China; Ruixinbio, Quanzhou, China) instructions.

Histology of Lung Tissue

After the sacrifice of rats on day 33, the left-lung lobes were removed rapidly and fixed in 4% paraformaldehyde for ≥ 24 h

at room temperature. Then, lung tissues were dehydrated in a graded series of ethanol solutions, cleared in xylene, and embedded in paraffin. Histology slices (thickness = 5 μ m) were prepared using a microtome. These slices were stained with hematoxylin and eosin (H&E) for morphology evaluation and inflammation using a microscope (BX50; Olympus, Tokyo, Japan) equipped with a digital camera (DFC 320, Leica, Wetzlar, Germany).

DNA Extraction and 16S rRNA Gene Sequencing of Bacteria

Thirty-five fresh feces from rats (seven from Con, five from Model, seven from Probio, eight from Hfat, and eight from Hfiber groups) were harvested aseptically with 2-mL sterilized Eppendorf tubes, respectively, and stored at -80°C immediately.

DNA isolation and 16S rRNA gene sequencing were provided by Novogene (Beijing, China), as described previously (Wang et al., 2016; Zhang et al., 2018), to ascertain the composition of the bacteria in the feces of rats in different groups. DNA was extracted using the cetyl trimethyl ammonium bromide/sodium dodecyl sulfonate (CTAB/SDS) method. The V3-V4 sequences of 16S rRNA genes were amplified by polymerase chain reaction (PCR) using specific primers (341 forward: 5'-CCTAYGGGRBGCASCAG-3'; 806 reverse: 5'-GGACTACNNGGTATCTAAT-3'). The PCR-amplified product was purified with a gel extraction kit (Qiagen, Hilden, Germany). After quantification, amplicons were normalized to generate a sequencing library and sequenced on a NovaSeq™ platform (Illumina, San Diego, CA, United States).

Alpha diversity, including Chao1 and ACE, was analyzed with Quantitative Insights Into Microbial Ecology (QIIME) 1.7.0¹. Beta diversity was ascertained using the non-metric multi-dimensional scaling (NMDS) of QIIME 1.7.0 to determine the similarity of species diversity in different samples. Statistical

¹www.qiime.org

significance was evaluated according to stress. Significant differences in the relative abundance of genera among strains were obtained by linear discriminant analysis (LDA) effect size (LEfSe).

Statistical Analyses

Continuous variables with a normal distribution were represented as the mean \pm SEM, and comparisons between groups were determined by the parametric Student's *t*-test. For parameters with a non-normal distribution, median and interquartile range (IQR) values are provided, and males were compared with females using the non-parametric Wilcoxon rank-sum test. Absolute and relative frequencies were calculated for qualitative data, and tested by the chi-square test or Fisher's exact test. Pearson's correlation was computed between intestinal microbiota and cytokines. $p < 0.05$ was considered significant. Missing values were not replaced and did not contribute to the analysis of the variable. Experiments were carried out at least twice. Data were analyzed using Prism 8.0 (GraphPad, San Diego, CA, United States).

Accession Number

The sequence data from this study are deposited in the GenBank Sequence Read Archive with the accession number SUB9930073.

RESULTS

Collection of Dietary Data

The data [age, body mass index (BMI), allergy history, and daily food intake] of 85 study participants [40 (47.1%) men and 45 (52.9%) women] are shown in **Table 1**. The median (IQR) age at study inclusion was 35 (27–48) years in men and 34 (26–43) years in women. Forty-nine (57.6%) participants reported a normal BMI (18.5–23.9 kg/m²).

Table 2 shows the percentages of major-diet components in study participants. A significant difference was registered in the intake of vegetables, soybeans, animal-based food, and condiments. Excessive intake of soybeans (58.18 \pm 49.45 vs. 39.34 \pm 34.41 g), animal-based food (meat: 220.77 \pm 117.58 vs. 108.51 \pm 103.15 g; fish and shrimps: 103.75 \pm 73.86 vs. 62.61 \pm 33.88 g; eggs: 47.20 \pm 29.13 vs. 34.20 \pm 25.88 g) and condiments (oil: 38.27 \pm 16.29 vs. 23.85 \pm 9.36 g; salt: 7.86 \pm 2.95 vs. 5.20 \pm 1.79 g) was common in people suffering from allergy compared with that in participants not suffering from allergy. An inadequate intake of vegetables was common in people suffering from allergy (319.43 \pm 125.15 g) compared with that in participants not suffering from allergy (387.83 \pm 124.62 g) (**Table 2**).

Different Diet Influences the Effect of *L. paracasei* on Allergic Symptoms

To evaluate allergy symptoms in rats, we measured the frequency of nasal rubs and sneezes per rat during the 15 min observation period after the last OVA challenge immediately. Rats in the Model group showed increased nasal rubs (vs. Con group, $p < 0.001$). With probiotics modulation, the number of nasal rubs decreased (vs. Model group, $p < 0.01$). High-fat diet intervention caused a significantly higher number of nasal rubs in probiotics-modulation AA rats ($p < 0.001$), compared with that in the Probio group, along with a reduced nasal rubs in Hfiber group but not significantly (**Figures 2A,B**). Whereas, the allergic symptoms of rats in Hfiber group was inhibited significantly (vs. Model group, $p < 0.001$). The incidence of sneezes in rats in all the groups maintained no significance. These results indicated that a high-fiber diet acted synergistically with *L. paracasei* intervention to alleviate allergic symptoms.

The level of OVA-specific IgE in serum was investigated to elucidate the effect of different diets on *L. paracasei* that modulated the allergic response. The serum OVA-specific IgE level was higher in the Model group than in the Con group


TABLE 1 | Characteristics of study participants.

	All (n = 85)		Male (n = 40)		Female (n = 45)		P _{for sex}
	n	%	n	%	n	%	
Age (years)							
18–26	21	24.7	9	42.8	12	57.2	0.532
27–35	27	31.8	13	48.1	14	51.9	0.723
36–44	14	16.5	6	42.9	8	57.1	0.365
45–53	16	18.8	8	50.0	8	50.0	0.332
54–62	7	8.2	4	57.1	3	42.9	0.246
BMI (kg/m²)							
Underweight (<18.5)	7	8.2	3	42.9	4	57.1	0.691
Normal (18.5–23.9)	49	57.6	15	30.6	34	69.4	0.670
Overweight (24–29.9)	28	32.9	21	75.0	7	25.0	0.417
Obesity (≥ 30)	1	1.2	1	100.0	0	0.0	—
Allergic							
Yes	44	51.8	19	43.2	25	56.8	0.518
No	41	48.2	21	51.2	20	48.8	

TABLE 2 | Components of food intake and its percentage in study participants.

Components		Percentage (%)										P value
Cereals	Intake amount (g)	<100	100–175	175–250	250–325	325–400	400–475	475–550	550–625	≥625		
	Cereals (%)	allergic	0.0	15.9	2.3	29.5	27.3	2.3	13.6	6.8	2.3	0.918
	non-allergic	0.0	4.9	0.0	36.6	29.3	14.6	7.3	2.4	4.9		
Vegetables	Intake amount (g)	<100	100–200	200–300	300–400	400–500	500–600	≥600				
	Vegetables (%)	allergic	2.3	18.2	6.8	27.3	38.6	6.8	0.0			0.014
	non-allergic	2.4	4.9	4.9	29.3	36.6	17.1	4.9				
Fruits	Intake amount (g)	<100	100–200	200–300	300–400	400–500	500–600	≥600				
	Fruits (%)	allergic	13.6	13.6	34.1	25.0	9.1	4.5	0.0			0.779
	non-allergic	7.3	17.1	39.0	29.8	2.4	4.9	2.4				
Soybeans	Intake amount (g)	<20	20–40	40–60	60–80	≥80						
	Soybeans (%)	allergic	11.4	15.9	34.1	29.5	9.1					0.046
	non-allergic	9.8	50.0	25.0	0.0	9.1						
Animals food	Intake amount (g)	<50	50–100	100–150	150–200	200–250	250–300	300–350	350–400	≥400		
	Meat (%)	allergic	4.5	4.5	27.3	0.0	15.9	25.0	11.4	4.5	6.8	<0.001
	non-allergic	14.6	53.7	12.2	0.0	4.9	4.9	2.4	2.4	4.9		
	Intake amount (g)	<50	50–100	100–150	150–200	≥200						
	Fish and shrimp (%)	allergic	25.0	15.9	22.7	25.0	11.4					0.002
	non-allergic	22.0	58.5	19.5	0.0	0.0						
Egg (%)	Intake amount (g)	<25	25–50	50–75	≥75							
	allergic	18.2	20.5	52.3	9.1							0.033
	non-allergic	19.5	68.3	4.9	7.3							
Condiments	Intake amount (g)	<25	25–50	≥50								
	Oil (%)	allergic	11.4	52.3	36.4							<0.001
	non-allergic	46.3	48.8	4.9								
	Intake amount (g)	<6	6–12	≥12								
Salt (%)	allergic	4.5	88.6	6.8							<0.001	
non-allergic	58.5	41.5	0.0									

Forty-four out of 85 individuals had suffered from allergy. Significant p-values are indicated in bold font.



($p < 0.01$). There was a trend toward reducing OVA-specific IgE with high-fiber diet treatment ($p = 0.063$), although not significantly (**Figure 2C**).

We counted the number of inflammatory cells obtained from BALF 24 h after the final challenge with OVA (**Figure 2D**). An increased total number of white blood cells (WBCs) in BALF was detected in the Model group compared with that in the Con group ($p < 0.001$). Level of WBCs after administered *L. paracasei* was lower in Probio group, although this difference was not statistically significant. However, when the rats consumed the high-fiber diet, the total number of inflammatory cells reduced markedly compared with that in the Model group ($p < 0.05$). These data suggested that the total number of WBCs in BALF was inhibited significantly by dietary consumption of fiber, whereas a high-fat diet aggravated the inflammatory response of AA.

Airway Inflammation in *L. paracasei*-Modulating AA Rats Fed Different Diets

To examine altered inflammation in allergic airways, histological examination of the lung tissue of AA rats was undertaken. Eosinophilic cytoplasm stained red with H&E (**Figure 3**). The typical pathological features of AA were observed in Model group as compared to Con group, along with discernible damage, edema, and a thickened mucosa. Probiotics treatment was effective in reducing the inflammation. Compared with the Model group, a more robust inflammatory response occurred in the Hfat group: extensive infiltration by eosinophils in peribronchial and alveolar septa, along with markedly thickened alveolar and bronchial walls. However, infiltration of inflammatory cells was suppressed significantly in the Hfiber group.

Alteration of Cytokine Levels From T Cells by Consumption of Different Diets Influenced *L. paracasei* Intervention

We measured cytokine levels in BALF and serum after the OVA challenge to further investigate the skewing of Th1/Th2 and Th17/T_{regs} immune responses after consuming different diets and how this influenced *L. paracasei* intervention. As shown in **Figure 4**, discrepancy is apparent in applying the *L. paracasei* modulating AA when there are different diets intervened. Notably, compared with the Hfat group, IL-4 was dropped

significantly in Hfiber group ($p < 0.05$). The Hfiber group contained apparently lower Th17 cytokine IL-17 level in serum compared with the Model group ($p < 0.05$). Increased production of IL-17 was observed intervened with high-fat diet, compared with Hfiber group ($p < 0.05$). Th1 cytokine IFN- γ in both BALF and serum were on the trend opposite from IL-4, although the differences were not statistical significant. The ratio of IL-4/IFN- γ was significantly different in Hfat group from that in Hfiber group ($P < 0.05$) (see **Supplementary Figure 1**). Based on these results, it was indicated that different diets affect the profile of cytokines in AA during the modulation of probiotics. The high-fiber diet blockaded the allergic inflammation significantly by regulating the levels of multiple inflammatory mediators therein, while the high-fat diet exacerbated it.

GM Composition of AA Rats Using 16S rRNA Gene Sequencing

A total of 1,915 operational taxonomic units (OTUs) were shared out of sequences obtained from 35 fecal samples. Nineteen phyla, 240 genera, and 162 species of gut microbes were annotated for subsequent analyses.

We wished to evaluate changes in the structure of the microbiota community among groups fed a different diet. Hence, microbial alpha diversity was measured using ACE and Chao1. As a result, we discovered a significantly higher alpha diversity for the Hfat group compared with that in the Probio group ($p < 0.01$) (**Figures 5A,B**). Furthermore, ordination with NMDS showed distinct spatial clustering between five groups (stress = 0.142) (**Figure 5C**).

To further illustrate the differences in gut microbiota composition among groups, we applied the LEfSe method to discover biomarkers in high-dimensional data and reveal genomic characteristics (**Figure 6**). Differentially enriched bacterial colonizers with LDA > 4 were identified among the five groups. Phylum Actinobacteria, families Bifidobacteriaceae and Peptostreptococcaceae, genera *Bifidobacterium*, *Turicibacter*, and *Romboutsia* were enriched in Con group. Family Erysipelotrichaceae and genus *Dubosiella* were enriched in Model group. Abundances of family Lactobacillaceae and genus *Lactobacillus* were enriched in Probio group. Abundances of phylum Firmicutes and Proteobacteria, families Lachnospiraceae, Ruminococcaceae and Desulfovibrionaceae, genera *Blautia*, *unidentified_Ruminococcaceae*, *unidentified_Clostridiales* and




FIGURE 3 | Histology of H&E-stained lung tissue (original magnification: $\times 100$). Photomicrographs were captured to detect eosinophil infiltration. Eosinophil infiltration was significantly higher in the Hfat group than the other groups. Interestingly, eosinophil inflammation was suppressed dramatically consumption of a high-fiber diet. $n = 5-8$ rat/group.




FIGURE 4 | Altered levels of the cytokines after OVA challenge. (A) TGF- β in BALF. (B) IL-17 in BALF. (C) IL-4 in BALF. (D) IFN- γ in BALF. (E) TGF- β in serum. (F) IL-17 in serum. (G) IL-4 in serum. (H) IFN- γ in serum. Data are the mean secretion (pg/mL) \pm SEM, $n = 4$; * $p < 0.05$, ** $p < 0.001$.





FIGURE 5 | Diversity in microbiota-community structure in different rat groups. (A) ACE indice. (B) Chao1 indice. * $p < 0.05$, ** $p < 0.01$. (C) NMDS. Stress < 0.2 , $\rho < 0.05$.

Oscillibacter were enriched in Hfat group. Biomarkers in Hfiber group including phylum Bacteroidetes, families Muribaculaceae, Tannerellaceae, Prevotellaceae, Enterococcaceae, genera *Allobaculum*, *Parabacteroides*, and *Enterococcus*.

To determine whether specific bacteria were differentially abundant along with the probiotics modulation intervened by different diets, we conducted a further analyses. After the AA model had been established, family Erysipelotrichaceae

and genera *Dubosiella* and *Paraprevotella* were more abundant, compared with that in the Con group (see **Supplementary Figure 2**). After intervention with *L. paracasei*, the dominant bacteria changed into phylum Bacteroidetes, and there were significant decrease in family Erysipelotrichaceae and genus *Dubosiella* (see **Supplementary Figure 3**). Upon consumption of different diets, the intestinal community altered significantly. In the Hfat group, phylum Proteobacteria,



families unidentified_Clostridiales, Desulfovibrionaceae, Ruminococcaceae and Lachnospiraceae, and genera *Anaerotruncus*, *Oscillibacter*, *Candidatus_Soleaferrea* and *Blautia* were dominant, while phylum Actinobacteria, families Lactobacillaceae, Bifidobacteriaceae and Muribaculaceae, genera *Lactobacillus*, *Bifidobacterium*, and *Turicibacter* decreased significantly (see **Supplementary Figure 4**). In the Hfiber group, the phylum Bacteroidetes, families Tannerellaceae, Lachnospiraceae and Muribaculaceae, genera *Angelakisella*, *Parabacteroides*, and *Allobaculum* were in relatively high abundance, phyla Firmicutes and Actinobacteria, families Lactobacillaceae, Bifidobacteriaceae, and Corynebacteriaceae, genera *Lactobacillus*, *Bifidobacterium*, *Turicibacter*, and *Corynebacterium* decreased (see **Supplementary Figure 5**). Taken together, these data demonstrated that different diets promoted distinct microbial communities that influenced *L. paracasei*-modulated AA.

Correlation Analysis for the Intestinal Microbiota and Cytokine Expression in BALF

The Pearson correlation coefficients between the predominant intestinal microbiota and cytokines in BALF were displayed in **Table 3**. The BALF level of IL-4 was positively correlated with

the abundance of Lachnospiraceae, Deltaproteobacteria, Desulfovibrionaceae, and Clostridiales, whereas TGF-β was negatively correlated with the abundance of Deltaproteobacteria and Desulfovibrionaceae, respectively ($p < 0.05$). The BALF level of IFN-γ was negatively associated with Erysipelotrichales abundance ($r = -0.483$, $p = 0.031$). A strong positive association and high significance between the IL-17 and abundance of *Turicibacter*, Peptostreptococcaceae, and *Romboutsia*, respectively, were found ($p < 0.05$).

DISCUSSION

Diet affects the GM, and dietary habits can impact overall intestinal health (Nakaji et al., 2002; Gophna, 2011; Wu et al., 2011; Bernaud and Rodrigues, 2013; Trompette et al., 2014). However, little is known of the role of diet in probiotic-modulating AA. One might speculate an interplay of diet, intestinal microflora, and immune cells. The diet pattern could be critical in regulating the effect of probiotics given to people suffering from AA.

The dietary patterns of 85 individuals were obtained. We found that people suffering from AA had unbalanced diets and inadequate intake of vegetables, and excessive intake of

TABLE 3 | Pearson correlation coefficients between intestinal bacteria and cytokines in BALF.

Intestinal bacteria	IL-4(pg/mL)		IFN- γ (pg/mL)		IL-17(pg/mL)		TGF- β (pg/mL)	
	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
Ruminococcaceae	0.393	0.087	-0.021	0.930	0.124	0.602	-0.380	0.098
<i>Lactobacillus</i>	-0.232	0.326	0.415	0.069	0.071	0.765	0.062	0.795
<i>Blautia</i>	0.078	0.743	-0.157	0.508	-0.183	0.440	-0.113	0.637
Bifidobacteriaceae	-0.158	0.505	0.060	0.801	0.185	0.435	0.402	0.079
<i>Turicibacter</i>	0.224	0.343	-0.048	0.840	0.502	0.024	0.019	0.938
Erysipelotrichaceae	0.055	0.816	-0.483	0.816	-0.018	0.939	0.126	0.597
<i>Oscillibacter</i>	0.233	0.323	0.076	0.750	0.084	0.725	-0.205	0.386
Peptostreptococcaceae	0.279	0.233	-0.218	0.357	0.612	0.004	-0.020	0.934
<i>Romboutsia</i>	0.211	0.372	-0.127	0.595	0.632	0.003	0.091	0.701
Lachnospiraceae	0.470	0.037	-0.098	0.681	-0.088	0.712	-0.294	0.208
Erysipelotrichales	0.055	0.816	-0.483	0.031	-0.018	0.939	0.126	0.597
Deltaproteobacteria	0.465	0.039	-0.062	0.794	-0.034	0.887	-0.527	0.017
<i>Bifidobacterium</i>	-0.158	0.505	0.060	0.801	0.185	0.435	0.402	0.079
Enterococcaceae	-0.097	0.684	-0.175	0.461	-0.273	0.244	-0.322	0.166
<i>Parabacteroides</i>	-0.325	0.162	0.090	0.705	-0.193	0.414	0.194	0.414
Prevotellaceae	-0.225	0.339	-0.005	0.983	-0.216	0.360	-0.009	0.971
Clostridiales	0.513	0.021	-0.115	0.629	0.070	0.768	-0.394	0.086
Tannerellaceae	-0.325	0.162	0.090	0.705	-0.193	0.414	0.194	0.414
<i>Allobaculum</i>	-0.254	0.279	-0.188	0.428	-0.390	0.089	0.229	0.332
Muribaculaceae	-0.334	0.150	0.031	0.897	-0.131	0.583	0.252	0.283
<i>Dubosiella</i>	-0.010	0.966	-0.377	0.101	-0.061	0.799	0.081	0.735
Lactobacillaceae	-0.232	0.326	0.415	0.069	0.071	0.765	0.062	0.795
Desulfovibrionaceae	0.478	0.033	-0.050	0.835	-0.026	0.913	-0.538	0.014
<i>Enterococcus</i>	-0.097	0.684	-0.175	0.461	-0.273	0.244	-0.322	0.166
<i>unidentified_Clostridiales</i>	0.425	0.061	-0.275	0.240	-0.081	0.732	-0.583	0.007

* $p < 0.05$, ** $p < 0.01$. $P < 0.05$ is highlighted in bold.

animal-based food was typical. According to a World Health Report from 2014, low intake of vegetables and fruit is associated with ischemic heart disease, gastrointestinal cancer, and stroke (World Health Organization (WHO), 2014). Consistent with the notion that a plant-rich diet is associated with positive health outcomes and reduced risk of disease, Alison et al. highlighted the importance of a high-fiber diet in protection against asthma development (Thorburn et al., 2015). Saadeh et al. (2015) observed that consumption of fast food and butter intake were associated with an increased prevalence of asthma symptoms among atopic children. A high-fat diet has been shown to increase airway inflammation in asthma (Wood et al., 2011). Hence, diet could be an environmental factor that influences how probiotics prevent AA.

We monitored systemic symptoms (nasal rubbing and sneezing), cytokines levels, inflammatory cell infiltration, and fecal bacterial composition in a dietary-intervention study of AA (induced by OVA) rats with modulation by *L. paracasei*. We discovered that different diets influenced the effect of *L. paracasei* on modulating AA. Moreover, a high fiber diet plus *L. paracasei* as a rescue medication showed a synergistic effect and led to a decreased frequency of rubbing and sneezes, whereas a high-fat diet exacerbated these symptoms. A reduced serum level of OVA-specific

IgE and total counts of inflammatory cells in BALF in the Hfiber group suggested an instinctive response to remission of allergic inflammation, but the group that consumed a high-fat diet showed the opposite results. IgE on the surface of immune cells binds to specific airway allergens followed by IgE cross-linking, cell activation, and release of preformed mediators (Komi and Bjermer, 2019; Salomonsson et al., 2019; Liu et al., 2020). In this regard, the lower IgE level also reflected attenuation of the Th2 immune response and the immune tolerance to allergic antigens (Ciprandi et al., 2015; Strokes and Casale, 2015). AA is characterized by eosinophils accumulation. The decreased number of eosinophils infiltrating the lungs was a response to the inhibition of allergic inflammation by consumption of a high-fiber diet. A high-fat diet induced eosinophil accumulation in the lungs of AA rats.

Th2 cells have crucial roles in AA pathogenesis, and imbalance of Th1/Th2 cells and disturbed balance of Th17/T_{reg} cells has been reported in asthma patients. Hence, to discriminate the state of Th1/Th2 and Th17/T_{reg} subsets in allergic airways, we measured the levels of IFN- γ (Th1 cytokine), IL-4 (Th2 cytokine), IL-17 (Th17 cytokine), and TGF- β (T_{reg} cytokine) by ELISA. The Probio group given a high-fiber diet inhibited expression of IL-4 and IL-17 and increased expression of

IFN- γ , but the group that consumed a high-fat diet had the opposite results. Th2 cytokines are believed to regulate IgE synthesis, and eosinophil numbers/activity are thought to play a significant part in driving AA pathogenesis (Dent et al., 1990; Savelkoul et al., 1991). IL-4 can upregulate expression of chemotactic factors such as eotaxins to promote eosinophil infiltration. IL-17 can drive airway remodeling (Quan et al., 2012). Indeed, our study found that the Hfiber diet assists the *L. paracasei* skewed the immune balance toward Th1, with increased IFN- γ and conversely decreased IL-4, in commitment with the previous presentation of reduced IgE and fewer eosinophils infiltration. Alteration in the balance of Th1/Th2 cytokines is an essential indicator of functional changes in suppressing the aberrant immune response in allergic diseases (Skapenko and Schulze-Koops, 2007; Cho et al., 2009). Although increased expression of IFN- γ has been found in individuals with severe asthma and acute exacerbations (Kumar et al., 2006; Barnes, 2009), there is evidence that probiotics promote IFN- γ production to reduce allergic inflammation (Giudice et al., 2010). Higher expression of IFN- γ in the Hfiber group with decreased expression of IL-4 compared with that in the Hfat group, and the ratio of IL-4/IFN- γ implied skewing of the balance of Th1/Th2 cells. The balance between Th17 and T_{regs} cells also suggests a crucial role in asthma pathogenesis (Vroman et al., 2018; Wang et al., 2018; Xu et al., 2018). Various studies have indicated that the homeostatic balance between T_{regs} and Th17 cells was altered markedly in asthma exacerbations, and correlates with asthma severity (Zou et al., 2018). TGF- β is produced by T_{regs}, which have been implicated to inhibit the immune response. They suppress effector T cells of Th1 or Th2 phenotypes. Therefore, increased expression of TGF- β in BALF in the Hfiber group seems rational.

Several studies have associated a changed intestinal microbiota with the etiology of various diseases, and changes in gut microflora composition, in response to diet change (Kau et al., 2011; David et al., 2014). We discovered that consumption of different diet led to alterations in the structure of microbial communities in the gut as well as the composition of intestinal bacteria as evidenced by NMDS (Gilotteaus et al., 2016). There was a significant difference in the alpha diversity of microbiota in different diet groups, and the bacterial diversity of the Hfiber group was decreased dramatically.

We revealed that a diet rich in fiber assisted the way that *L. paracasei* modified the GM. This finding corroborates the data from a study by Salonen et al. (2014), who reported that a diet high in fiber was associated with decreased bacterial diversity relative to that of other diets. Comparison of 16S rRNA gene sequencing among groups revealed that a high-fiber diet significantly improved GM structure by enriching bacteria of the phylum Bacteroidetes, as reported by Trompette et al. (2014). Consumption of a high-fat diet enhanced the proportions of Proteobacteria and Firmicutes, which indicated the enrichment of specific bacteria in the colon is diet-specific (Filippo et al., 2010; Walker et al., 2011; Campo et al., 2014; Cruz et al., 2014; Ramos-Romero

et al., 2018). Phylum Bacteroidetes was increased markedly in the Hfiber group. Salyers and coworkers showed that the bacteria of phylum Bacteroidetes have the functions of carbohydrate fermentation, polysaccharide metabolism, and maintaining the normal physiological function of the intestinal tract (Salyers, 1984). The influence of a high-fat diet led to significant induction of Clostridiales, Lachnospiraceae, Ruminococcaceae, Deltaproteobacteria, and Desulfococcaceae. Studies have shown that Clostridiales can be pathogenic, leading to infectious diseases or mild cognitive impairment in mice (Antharam et al., 2013; Vogt et al., 2017). The members of the family Lachnospiraceae are enriched by a low-fiber Western-style diet, are associated with inflammatory diseases, and have been reported to protect against allergy by modulating the immune system (Png et al., 2010; Pascal et al., 2018). The relative abundance of bacteria of the family Ruminococcaceae is lower in IgE-associated eczema compared with that in people not suffering from allergies. Enriched abundance of Deltaproteobacteria is related to AA in mice (Hirota et al., 2019). Moreover, bacteria of the family Desulfococcaceae (which are opportunistic pathogens) have been linked to inflammatory diseases of the gut and chronic disorders. Hence, the alterations induced by a high-fat diet suggest inflammation exacerbation.

Our correlation analysis of intestinal bacteria and cytokine expression (section “Correlation Analysis for the Intestinal Microbiota and Cytokine Expression in BALF”) demonstrated that GM alterations might be associated with inflammation during AA. Therefore, *L. paracasei*, with the assistance of a high-fiber diet, ameliorated allergy symptoms to a greater extent than use of *L. paracasei* alone in an inflammatory-modulation manner. Thus, a high fiber diet seems to support the effect of AA medication as an add-on therapy.

Our study had two main limitations. First, our findings are limited to an AA model induced by OVA; the effect a high-fiber or high-fat diet on an AA sensitized by other antigens (e.g., house dust mite) is not known. Second, associations between the most relevant taxa and a high-fiber diet or high-fat diet, respectively, were not validated by real-time reverse transcription-quantitative PCR.

CONCLUSION

We provided new insights into consumption of different diets influenced the effect of *L. paracasei* on suppressing AA. In particular, the intestinal microbiota altered by different dietary patterns were associated with allergic inflammation.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories

and accession number(s) can be found in the article/**Supplementary Material**.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Ethics Committee of Dalian Medical University. The patients/participants provided their written informed consent to participate in this study. The animal study was reviewed and approved by the Animal Ethics Committee of Dalian Medical University.

AUTHOR CONTRIBUTIONS

SW conceived and designed the experiments and supervised. AX, JS, and SL performed the experiments. JS and YL analyzed the data. AX wrote the manuscript. SW and LT reviewed the manuscript. All authors have read and given approval to the final version of the manuscript.

REFERENCES

- Abrahamsson, T. R., Jakobsson, H. E., Andersson, A. F., Björkstén, B., Engstrand, L., and Jenmalm, M. C. (2014). Low gut microbiota diversity in early infancy precedes asthma at school age. *Clin. Exp. Allergy* 44, 842–850.
- Antharam, V. C., Li, E. C., Ishmael, A., Sharma, A., Mai, V., Rand, K. H., et al. (2013). Intestinal dysbiosis and depletion of butyrogenic bacteria in *Clostridium difficile* infection and nosocomial diarrhea. *J. Clin. Microbiol.* 51, 2884–2892. doi: 10.1128/jcm.00845-13
- Atarashi, K., Tanoue, T., Shima, T., Imaoka, A., Kuwahara, T., Momose, Y., et al. (2011). Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science* 331, 337–341. doi: 10.1126/science.1198469
- Barnes, P. J. (2009). The cytokine network in chronic obstructive pulmonary disease. *Am. J. Respir. Cell Mol. Biol.* 41, 631–638. doi: 10.1165/rcmb.2009-0220tr
- Bernaudo, F. S. R., and Rodrigues, T. C. (2013). Dietary fiber – adequate intake and effects on metabolism health. *Arq. Bras. Endocrinol. Metabol.* 57, 397–405. doi: 10.1590/s0004-27302013000600001
- Campo, R., Garriga, M., Pérez-Aragón, A., Guallarte, P., Lamas, A., Máiz, L., et al. (2014). Improvement of digestive health and reduction in proteobacterial populations in the gut microbiota of cystic fibrosis patients using a *Lactobacillus reuteri* probiotic preparation: a double blind prospective study. *J. Cyst. Fibros.* 13, 716–722. doi: 10.1016/j.jcf.2014.02.007
- Cho, K.-S., Park, H.-K., Park, H.-Y., Jung, J. S., Jeon, S.-G., Kim, Y.-K., et al. (2009). IFATS collection: immunomodulatory effects of adipose tissue-derived stem cells in an allergic rhinitis mouse model. *Stem Cells* 27, 259–265. doi: 10.1634/stemcells.2008-0283
- Choi, S.-P., Oh, H.-N., Choi, C.-Y., Ahn, H., Yun, H. S., Chung, Y. M., et al. (2018). Oral administration of *Lactobacillus plantarum* CJLP133 and CJLP243 alleviates birch pollen-induced allergic rhinitis in mice. *J. Appl. Microbiol.* 124, 821–828. doi: 10.1111/jam.13635
- Ciprandi, G., Marseglia, G. L., Castagnoli, R., Valsecchi, C., Tagliacarne, C., Caimmi, S., et al. (2015). From IgE to clinical trials of allergic rhinitis. *Expert Rev. Clin. Immunol.* 11, 1321–1333.
- Colombo, D., Zagni, E., Ferri, F., Canonica, G. W., and Proxima study centers. (2019). Gender differences in asthma perception and its impact on quality of life: a post hoc analysis of the PROXIMA (Patient Reported Outcomes and Xolair® In the Management of Asthma) study. *Allergy Asthma Clin. Immunol.* 15:65. doi: 10.1186/s13223-019-0380-z

FUNDING

This work was funded by National Natural Science Foundation of China (No. 81370113).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.737622/full#supplementary-material>

Supplementary Figure 1 | The ratio of IL-4/IFN- γ .

Supplementary Figure 2 | Gut microbiota comparisons between Con group and Model group.

Supplementary Figure 3 | Gut microbiota comparisons between Model group and Probio group.

Supplementary Figure 4 | Gut microbiota comparisons between Probio group and Hfat group.

Supplementary Figure 5 | Gut microbiota comparisons between Probio group and Hfiber group.

Supplementary Table 1 | Diet composition.

- Cruz, P. D., Kang, S., Wagner, J., Buckley, M., Sim, W. H., Prideaux, L., et al. (2014). Specific mucosa-associated microbiota in Crohn's disease at the time of resection are associated with early disease recurrence: a Pilot Study. *J. Gastroenterol. Hepatol.* 30, 268–278. doi: 10.1111/jgh.12694
- David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., et al. (2014). Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 505, 559–563. doi: 10.1038/nature12820
- Dent, L. A., Strath, M., Mellor, A. L., and Sanderson, C. J. (1990). Eosinophilia in transgenic mice expressing interleukin 5. *J. Exp. Med.* 172, 1425–1431. doi: 10.1084/jem.172.5.1425
- Dzidic, M., Abrahamsson, T. R., Artacho, A., Björkstén, B., Collado, M. C., Mira, A., et al. (2017). Aberrant IgA responses to the gut microbiota during infancy precede asthma and allergy development. *J. Allergy Clin. Immunol.* 139, 1017–1025. doi: 10.1016/j.jaci.2016.06.047
- Ellwood, P., Asher, M. I., Björkstén, B., Burr, M., Pearce, N., and Robertson, C. F. (2001). Diet and asthma, allergic rhinoconjunctivitis and atopic eczema symptom prevalence: an ecological analysis of the International Study of Asthma and Allergies in Childhood (ISAAC) data. *Eur. Respir. J.* 17, 436–443. doi: 10.1183/09031936.01.17304360
- Filippo, C. D., Cavalieri, D., Paola, M. D., Ramazzotti, M., Poullet, J. B., Massart, S., et al. (2010). Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc. Natl. Acad. Sci. U. S. A.* 107, 14691–14696. doi: 10.1073/pnas.1005963107
- Fiocchi, A., Burks, W., Bahna, S. L., Bielory, L., Boyle, R. J., Cocco, R., et al. (2012). Clinical use of probiotics in pediatric allergy (CUPPA): a World Allergy Organization position paper. *World Allergy Organ J.* 5, 148–167. doi: 10.1097/wox.0b013e3182784ee0
- Frei, R., Lauener, R. P., Cramer, R., and O'Mahony, L. (2012). Microbiota and dietary interactions: an update to the hygiene hypothesis? *Allergy* 67, 451–461. doi: 10.1111/j.1398-9995.2011.02783.x
- Fujiwara, D., Inoue, S., Wakabayashi, H., and Fujii, T. (2004). The anti-allergic effects of lactic acid bacteria are strain dependent and mediated by effects on both Th1/Th2 cytokine expression and balance. *Int. Arch. Allergy Immunol.* 135, 205–215. doi: 10.1159/000081305
- Giloteaux, L., Goodrich, J. K., Walters, W. A., Levine, S. M., Ley, R. E., and Hanson, M. R. (2016). Reduced diversity and altered composition of the gut microbiome in individuals with myalgic encephalomyelitis/chronic fatigue syndrome. *Microbiome* 4:30. doi: 10.1186/s40168-016-0171-4

- Giudice, M. M., Leonardi, S., Maiello, N., and Brunese, F. P. (2010). Food allergy and probiotics in childhood. *J. Clin. Gastroenterol.* 44, S22–S25. doi: 10.1097/mcg.0b013e3181e102a7
- Gollwitzer, E. S., Saglani, S., Trompette, A., Yadava, K., Sherburn, R., McCoy, K. D., et al. (2014). Lung microbiota promotes tolerance to allergens in neonates via PD-L1. *Nat. Med.* 20, 642–647. doi: 10.1038/nm.3568
- Gophna, U. (2011). Microbiology. The guts of dietary habits. *Science* 334, 45–46. doi: 10.1126/science.1213799
- Grimshaw, K. E. C., Maskell, J., Oliver, E. M., Morris, R. C. G., Foote, K. D., Mills, E. N. C., et al. (2014). Diet and food allergy development during infancy: birth cohort study findings using prospective food diary data. *J. Allergy Clin. Immunol.* 133, 511–519. doi: 10.1016/j.jaci.2013.05.035
- Herbst, T., Sichelstiel, A., Schär, C., Yadava, K., Bürki, K., Cahenzli, J., et al. (2011). Dysregulation of allergic airway inflammation in the absence of microbial colonization. *Am. J. Respir. Crit. Care Med.* 184, 198–205. doi: 10.1164/rccm.201010-1574OC
- Hirota, R., Ohya, Y., Yamamoto-Hanada, K., Fukutomi, Y., Muto, G., Ngatu, N. R., et al. (2019). Triclosan-induced alteration of gut microbiome and aggravation of asthmatic airway response in aeroallergen-sensitized mice. *Allergy* 74, 996–999. doi: 10.1111/all.13639
- Hou, J. K., Abraham, B., and El-Serag, H. (2011). Dietary intake and risk of developing inflammatory bowel disease: a systematic review of the literature. *Am. J. Gastroenterol.* 106, 563–573. doi: 10.1038/ajg.2011.44
- Jensen, M. P., Meldrum, S., Taylor, A. L., Dunstan, J. A., and Prescott, S. L. (2012). Early probiotic supplementation for allergy prevention: long-term outcomes. *J. Allergy Clin. Immunol.* 130, 1209–1211. doi: 10.1016/j.jaci.2012.07.018
- Julia, V., Macia, L., and Dombrowicz, D. (2015). The impact of diet on asthma and allergic diseases. *Nat. Rev. Immunol.* 15, 308–322. doi: 10.1038/nri3830
- Kau, A. L., Ahern, P. P., Griffin, N. W., Goodman, A. L., and Gordon, J. I. (2011). Human nutrition, the gut microbiome and the immune system. *Nature* 474, 327–336. doi: 10.1038/nature10213
- Komi, D. E. A., and Bjermer, L. (2019). Mast cell-mediated orchestration of the immune responses in human allergic asthma: current insights. *Clin. Rev. Allergy Immunol.* 56, 234–247. doi: 10.1007/s12016-018-8720-1
- Kuitunen, M., Kukkonen, K., Juntunen-Backman, K., Korpela, R., Poussa, T. T., Haahela, T., et al. (2009). Probiotics prevent IgE-associated allergy until age 5 years in cesarean-delivered children but not in the total cohort. *J. Allergy Clin. Immunol.* 123, 335–341. doi: 10.1016/j.jaci.2008.11.019
- Kumar, R. K., Webb, D. C., Herbert, C., and Foster, P. S. (2006). Interferon-gamma as a possible target in chronic asthma. *Inflamm. Allergy Drug Targets* 5, 253–256. doi: 10.2174/187152806779010909
- Lin, C.-H., Tseng, C.-Y., and Chao, M.-W. (2020). Administration of *Lactobacillus paracasei* HB89 mitigates PM_{2.5}-induced enhancement of inflammation and allergic airway response in murine asthma model. *PLoS One* 15:e0243062. doi: 10.1371/journal.pone.0243062
- Liu, H., Tan, J., Liu, J., Feng, H., and Pan, D. (2020). Altered mast cell activity in response to rhinovirus infection provides novel insight into asthma. *J. Asthma* 57, 459–467. doi: 10.1080/02770903.2019.1585870
- Makino, T., Yamashita, M., Takeuchi, N., Kabuki, T., Hattori, M., and Yoshida, T. (2019). *Lactobacillus helveticus* SBT2171 alleviates allergic symptoms in a murine model for pollen allergy. *Biosci. Biotechnol. Biochem.* 83, 2298–2306. doi: 10.1080/09168451.2019.1654847
- Manzel, A., Muller, D. N., Hafler, D. A., Erdman, S. E., Linker, R. A., and Kleinewietfeld, M. (2014). Role of “western diet” in inflammatory autoimmune diseases. *Curr. Allergy Asthma Rep.* 14:404. doi: 10.1007/s11882-013-0404-6
- Maslowski, K. M., Vieira, A. T., Ng, A., Kranich, J., Sierro, F., Yu, D., et al. (2009). Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 461, 1282–1286. doi: 10.1038/nature08530
- Myles, I. A. (2014). Fast food fever: reviewing the impacts of the Western diet on immunity. *Nutr. J.* 13:61. doi: 10.1186/1475-2891-13-61
- Nakaji, S., Sugawara, K., Saito, D., Yoshioka, Y., MacAuley, D., Bradley, T., et al. (2002). Trends in dietary fiber intake in Japan over the last century. *Eur. J. Nutr.* 41, 222–227. doi: 10.1007/s00394-002-0379-x
- Pascal, M., Perez-Gordo, M., Caballero, T., Escribese, M. M., Longo, M. N. L., Luengo, O., et al. (2018). Microbiome and allergic diseases. *Front. Immunol.* 9:1584. doi: 10.3389/fimmu.2018.01584
- Png, C. W., Lindén, S. K., Gilshenan, K. S., Zoetendal, E. G., McSweeney, C. S., Sly, L. I., et al. (2010). Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. *Am. J. Gastroenterol.* 105, 2420–2428. doi: 10.1038/ajg.2010.281
- Protudjer, J. L. P., Sevenhuysen, G. P., Ramsey, C. D., Kozyrskyj, A. L., and Becker, A. B. (2012). Low vegetable intake is associated with allergic asthma and moderate-to-severe airway hyperresponsiveness. *Pediatr. Pulmonol.* 47, 1159–1169. doi: 10.1002/ppul.22576
- Quan, S.-H., Zhang, Y.-L., Han, D. H., Iwakura, Y., and Rhee, C.-S. (2012). Contribution of interleukin 17A to the development and regulation of allergic inflammation in a murine allergic rhinitis model. *Ann. Allergy Asthma Immunol.* 108, 342–350. doi: 10.1016/j.anaai.2012.02.014
- Ramos-Romero, S., Hereu, M., Atienza, L., Casas, J., Jáuregui, O., Amézqueta, S., et al. (2018). Mechanistically different effects of fat and sugar on insulin resistance, hypertension, and gut microbiota in rats. *Am. J. Physiol. Endocrinol. Metab.* 314, E552–E563. doi: 10.1152/ajpendo.00323.2017
- Saadeh, D., Salameh, P., Caillaud, D., Charpin, D., Blay, F. D., Kopferschmitt, C., et al. (2015). Prevalence and association of asthma and allergic sensitization with dietary factors in schoolchildren: data from the french six cities study. *BMC Public Health* 15:993. doi: 10.1186/s12889-015-2320-2
- Salomonsson, M., Malinovsky, A., Kalm-Stephens, P., Dahlin, J. S., Janson, C., Alving, K., et al. (2019). Circulating mast cell progenitors correlate with reduced lung function in allergic asthma. *Clin. Exp. Allergy* 49, 874–882. doi: 10.1111/cea.13388
- Salonen, A., Lahti, L., Salojärvi, J., Holtrop, G., Korpela, K., Duncan, S. H., et al. (2014). Impact of diet and individual variation on intestinal microbiota composition and fermentation products in obese men. *ISME J.* 8, 2218–2230. doi: 10.1038/ismej.2014.63
- Salyers, A. A. (1984). *Bacteroides* of the human lower intestinal tract. *Annu. Rev. Microbiol.* 38, 293–313. doi: 10.1146/annurev.mi.38.100184.001453
- Savelkoul, H. F., Seymour, B. W., Sullivan, L., and Coffman, R. L. (1991). IL-4 can correct defective IgE production in SJA/9 mice. *J. Immunol.* 146, 1801–1805.
- Schabussova, I., Hufnagl, K., Tang, M. L. K., Hoflehner, E., Wagner, G. L., Nutten, S., et al. (2012). Perinatal maternal administration of *Lactobacillus paracasei* NCC 2461 prevents allergic inflammation in a mouse model of birch pollen allergy. *PLoS One* 7:e40271. doi: 10.1371/journal.pone.0040271
- Shinmei, Y., Yano, H., Kagawa, Y., Izawa, K., Akagi, M., Inoue, T., et al. (2009). Effect of Brazilian propolis on sneezing and nasal rubbing in experimental allergic rhinitis of mice. *Immunopharmacol. Immunotoxicol.* 31, 688–693. doi: 10.3109/08923970903078443
- Skapenko, A., and Schulze-Koops, H. (2007). Analysis of Th1/Th2 T-cell subsets. *Methods Mol. Med.* 136, 87–96. doi: 10.1007/978-1-59745-402-5_7
- Strokes, J. R., and Casale, T. B. (2015). The use of anti-IgE therapy beyond allergic asthma. *J. Allergy Clin. Immunol. Pract.* 3, 162–166. doi: 10.1016/j.jaip.2014.10.010
- Thorburn, A. N., Macia, L., and Mackay, C. R. (2014). Diet, metabolites, and “western-lifestyle” inflammatory diseases. *Immunity* 40, 833–842. doi: 10.1016/j.immuni.2014.05.014
- Thorburn, A. N., McKenzie, C. I., Shen, S., Stanley, D., Macia, L., Masson, L. J., et al. (2015). Evidence that asthma is a developmental origin disease influenced by maternal diet and bacterial metabolites. *Nat. Commun.* 6:7320. doi: 10.1038/ncomms8320
- Trompette, A., Gollwitzer, E. S., Yadava, K., Sichelstiel, A. K., Sprenger, N., Ngom-Bru, C., et al. (2014). Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat. Med.* 20, 159–166. doi: 10.1038/nm.3444
- Tsabouri, S., Priftis, K. N., Chaliasos, N., and Siamopoulou, A. (2014). Modulation of gut microbiota downregulates the development of food allergy in infancy. *Allergol. Immunopathol.* 42, 69–77. doi: 10.1016/j.aller.2013.03.010
- Vogt, N. M., Kerby, R. L., Harding, S., Merluzzi, A. P., Koenig, L., Beilfuss, M., et al. (2017). Gut microbiome alterations in Alzheimer’s disease and relationship with Csf Biomarkers. *Alzheimers Dement.* 13:563. doi: 10.1016/j.jalz.2017.07.172
- Vroman, H., Bergen, I. M., Hulst, J. A. C., Nimwegen, M., Uden, D., Schuijs, M. J., et al. (2018). TNF- α -induced protein 3 levels in lung dendritic cells instruct TH2 or TH17 cell differentiation in eosinophilic or neutrophilic asthma. *J. Allergy Clin. Immunol.* 141, 1620–1621. doi: 10.1016/j.jaci.2017.08.012

- Walker, A. W., Ince, J., Duncan, S. H., Webster, L. M., Holtrop, G., Ze, X., et al. (2011). Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *ISME J.* 5, 220–230. doi: 10.1038/ismej.2010.118
- Wang, L., Li, P., Tang, Z., Yan, X., and Feng, B. (2016). Structural modulation of the gut microbiota and the relationship with body weight: compared evaluation of liraglutide and saxagliptin treatment. *Sci. Rep.* 6:33251. doi: 10.1038/srep33251
- Wang, L., Wan, H., Tang, W., Ni, Y., Hou, X., Pan, L., et al. (2018). Critical roles of adenosine A2A receptor in regulating the balance of Treg/Th17 cells in allergic asthma. *Clin. Respir. J.* 12, 149–157. doi: 10.1111/crj.12503
- Wang, X., Hui, Y., Zhao, L., Hao, Y., and Guo, H. (2017). Oral administration of *Lactobacillus paracasei* L9 attenuates PM2.5-induced enhancement of airway hyperresponsiveness and allergic airway response in murine model of asthma. *PLoS One* 12:e0171721. doi: 10.1371/journal.pone.0171721
- West, C. E., Hammarström, M.-L., and Hernell, O. (2013). Probiotics in primary prevention of allergic disease - follow-up at 8-9 years of age. *Allergy* 68, 1015–1020. doi: 10.1111/all.12191
- Wood, L. G., Garg, M. L., and Gibson, P. G. (2011). A high-fat challenge increases airway inflammation and impairs bronchodilator recovery in asthma. *J. Allergy Clin. Immunol.* 127, 1133–1140. doi: 10.1016/j.jaci.2011.01.036
- World Health Organization (WHO). (2014). *Promoting Fruit and Vegetable Consumption Around the World*. Switzerland: WHO.
- Wu, G. D., Chen, J., Hoffmann, C., Birringer, K., Chen, Y.-Y., and Keilbaugh, S. A. (2011). Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334, 105–108. doi: 10.1126/science.1208344
- Xu, L., Sun, W. J., Jia, A. J., Qiu, L. L., Xiao, B., Mu, L., et al. (2018). MBD2 regulates differentiation and function of Th17 cells in neutrophils-dominant asthma via HIF-1 alpha. *J. Inflamm.* 15:15. doi: 10.1186/s12950-018-0191-x
- Zhang, B., Sun, W., Yu, N., Sun, J., Yu, X., Li, X., et al. (2018). Anti-diabetic effect of baicalein is associated with the modulation of gut microbiota in streptozotocin and high-fat-diet induced diabetic rats. *J. Funct. Foods* 46, 256–267. doi: 10.1016/j.jff.2018.04.070
- Zhang, Z., Shi, L., Pang, W., Liu, W., Li, J., Wang, H., et al. (2016). Dietary fiber intake regulates intestinal microflora and inhibits ovalbumin-induced allergic airway inflammation in a mouse model. *PLoS One* 11:e0147778. doi: 10.1371/journal.pone.0147778
- Zou, X.-L., Chen, Z.-G., Zhang, T.-T., Feng, D.-Y., Li, H.-T., and Yang, H.-L. (2018). Th17/Treg homeostasis, but not Th1/Th2 homeostasis, is implicated in exacerbation of human bronchial asthma. *Ther. Clin. Risk Manag.* 14, 1627–1636. doi: 10.2147/tcrm.s172262

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Xie, Song, Lu, Liu, Tang and Wen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.